

Multiple scattering of polarized light by particles in an absorbing host medium

Janna M. Dlugach^{a,*} and Michael I. Mishchenko^a

^a*Main Astronomical Observatory of the National Academy of Sciences of Ukraine, 27 Zabolotny Str., 03680, Kyiv, Ukraine*

^b*NASA Goddard Institute for Space Studies, 2880 Broadway, New York, NY 10025, USA*

**Presenting author (dl@mao.kiev.ua)*

We use recently developed first-principles tools [1–3] to solve the vector radiative transfer equation for a particulate layer imbedded in an absorbing host medium. The results of extensive computations for typical geophysical scenarios are analyzed, and conclusions are formulated as to the practical importance of the effects of absorption in the host medium.

References

- [1] Mishchenko, M. I., 2008: Multiple scattering by particles embedded in an absorbing medium. 2. Radiative transfer equation. *J. Quant. Spectrosc. Radiat. Transfer* **109**, 2386–2390.
- [2] Mishchenko, M. I., and P. Yang, 2018: Far-field Lorenz–Mie scattering in an absorbing host medium: theoretical formalism and FORTRAN program. *J. Quant. Spectrosc. Radiat. Transfer* **205**, 241–252.
- [3] Mishchenko, M. I., J. M. Dlugach, J. A. Lock, and M. A. Yurkin, 2018: Far-field Lorenz–Mie scattering in an absorbing host medium. II: Improved stability of the numerical algorithm. *J. Quant. Spectrosc. Radiat. Transfer* **217**, 274–277.

Preferred mode of presentation: Oral